

- Helmrich, S. P., Ragland, D. R., Leung, R. W., & Paffenbarger Jr, R. S. (1991). Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *New England journal of medicine*, 325(3), 147-152.
- Hobbs, F. D., Gensini, G., Mancini, G. B., Manolis, A. J., Bauer, B., Böehler, S., & Marques da Silva, P. (2006). Can combining different risk interventions into a single formulation contribute to improved cardiovascular disease risk reduction? Rationale and design for an international, open-label program to assess the effectiveness of a single pill (amlodipine/atorvastatin) to attain recommended target levels for blood pressure and lipids (The JEWEL Program). *International journal of cardiology*, 110(2), 242-250.
- Hoffart, E., Ghebregiorghis, L., Nussler, A. K., Thasler, W. E., Weiss, T. S., Schwab, M., & Burk, O. (2012). Effects of atorvastatin metabolites on induction of drug-metabolizing enzymes and membrane transporters through human pregnane X receptor. *British journal of pharmacology*, 165(5), 1595-1608.
- Holman, R. R., Paul, S. K., Bethel, M. A., Matthews, D. R., & Neil, H. A. W. (2008). 10-year follow-up of intensive glucose control in type 2 diabetes. *New England Journal of Medicine*, 359(15), 1577-1589.
- Holman, R. R., Paul, S. K., Bethel, M. A., Neil, H. A. W., & Matthews, D. R. (2008). Long-term follow-up after tight control of blood pressure in type 2 diabetes. *New England Journal of Medicine*, 359(15), 1565-1576.