

| | | |
|-------------|---|----|
| Figure 3.24 | Chromatogram of amlodipine, glimepiride, and atorvastatin when setting the temperature at 22°C | 61 |
| Figure 3.25 | Chromatogram of amlodipine, glimepiride, and atorvastatin when prepared in plasma and temperature set at 28°C | 62 |
| Figure 3.26 | Chromatogram of amlodipine, glimepiride, and atorvastatin when prepared in plasma and temperature set at 22°C | 63 |
| Figure 3.27 | Chromatogram of amlodipine, glimepiride, and atorvastatin assayed in 5% increase of ACN. | 64 |
| Figure 3.28 | Chromatogram of amlodipine, glimepiride, and atorvastatin assayed in 5% decrease of ACN. | 64 |
| Figure 3.29 | Chromatogram of amlodipine, glimepiride, and atorvastatin assayed in 5% increase of ACN in plasma | 65 |
| Figure 3.30 | Chromatogram of amlodipine, glimepiride, and atorvastatin assayed in 5% decrease of ACN in plasma | 66 |
| Figure 3.31 | Chromatogram of amlodipine, glimepiride, and atorvastatin following 4 hours of sunlight exposure | 66 |
| Figure 3.32 | Chromatogram of placebo excipients | 67 |
| Figure 3.33 | Chromatogram of placebo in plasma | 68 |