

## References

- ALLHAT, O., the, A., & Coordinators for the ALLHAT Collaborative Research Group. (2002). Major outcomes in moderately hypercholesterolemic, hypertensive patients randomized to pravastatin vs usual care: the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT-LLT). *JAMA: the journal of the American Medical Association*, 288(23), 2998.
- Alsarra, I. A. (2009). High-performance liquid chromatographic method for quantitative determination of amlodipine in human plasma and pharmaceutical dosage form and its application to pharmacokinetic studies. *Journal of chromatographic science*, 47(10), 863-867.
- Altuntas, T. G., & Erk, N. (2004). Liquid chromatographic determination of atorvastatin in bulk drug, tablets, and human plasma. *Journal of liquid chromatography & related technologies*, 27(1), 83-93.
- Arguello, C. R. Update on Type 2 Diabetes. *Hemoglobin*, 1, 7-0.
- Asia Pacific Cohort Studies Collaboration. (2003). Cholesterol, coronary heart disease, and stroke in the Asia Pacific region. *International journal of epidemiology*, 32(4), 563-572.
- Athyros, V. G., Mikhailidis, D. P., Papageorgiou, A. A., Symeonidis, A. N., Pehlivanidis, A. N., Bouloukos, V. I., & Elisaf, M. (2004). The effect of statins versus untreated dyslipidaemia on renal function in patients with coronary heart disease. A subgroup analysis of the Greek atorvastatin and coronary heart disease evaluation (GREACE) study. *Journal of clinical pathology*, 57(7), 728-734.