

- Deshpande, A. D., Harris-Hayes, M., & Schootman, M. (2008). Epidemiology of diabetes and diabetes-related complications. *Physical Therapy*, 88(11), 1254-1264.
- Devabhaktuni, M., & Bangalore, S. (2009). Fixed combination of amlodipine and atorvastatin in cardiovascular risk management: patient perspectives. *Vasc Health Risk Manag*, 5(1), 377-387.
- Dolor, R. J., Yancy, W. S., Owen, W. F., Matchar, D. B., Samsa, G. P., Pollak, K. I., & Svetkey, L. P. (2009). Hypertension Improvement Project (HIP): study protocol and implementation challenges. *Trials*, 10(1), 13.
- Downs, J. R., Clearfield, M., Weis, S., Whitney, E., Shapiro, D. R., Beere, P. A., & AFCAPS/TexCAPS Research Group. (1998). Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. *Jama*, 279(20), 1615-1622.
- Duckworth, W., Abraira, C., Moritz, T., Reda, D., Emanuele, N., Reaven, P. D., & Huang, G. D. (2009). Glucose control and vascular complications in veterans with type 2 diabetes. *New England Journal of Medicine*, 360(2), 129-139.
- Erdine, S., Ro, Y. M., Tse, H. F., Howes, L. G., Aguilar-Salinas, C. A., Chaves, H., & Schou, I. M. (2009). Single-pill amlodipine/atorvastatin helps patients of diverse ethnicity attain recommended goals for blood pressure and lipids (the Gemini-AALA study). *Journal of human hypertension*, 23(3), 196-210.