

- Brandle, M., Zhou, H., Smith, B. R., Marriott, D., Burke, R., Tabaei, B. P., & Herman, W. H. (2003). The direct medical cost of type 2 diabetes. *Diabetes care*, 26(8), 2300-2304.
- Bremer, J. P., Jauch-Chara, K., Hallschmid, M., Schmid, S., & Schultes, B. (2009). Hypoglycemia unawareness in older compared with middle-aged patients with type 2 diabetes. *Diabetes care*, 32(8), 1513-1517.
- Brunham, L. R., Lansberg, P. J., Zhang, L., Miao, F., Carter, C., Hovingh, G. K., & Hayden, M. R. (2011). Differential effect of the rs4149056 variant in *SLCO1B1* on myopathy associated with simvastatin and atorvastatin. *The pharmacogenomics journal*, 12(3), 233-237.
- Buchwald, H., Varco, R. L., Matts, J. P., Long, J. M., Fitch, L. L., Campbell, G. S., & Chalmers, T. C. (1990). Effect of partial ileal bypass surgery on mortality and morbidity from coronary heart disease in patients with hypercholesterolemia: report of the Program on the Surgical Control of the Hyperlipidemias (POSCH). *New England Journal of Medicine*, 323(14), 946-955.
- Chrysant, S. G., Melino, M., Karki, S., Lee, J., & Heyrman, R. (2008). The combination of olmesartan medoxomil and amlodipine besylate in controlling high blood pressure: COACH, a randomized, double-blind, placebo-controlled, 8-week factorial efficacy and safety study. *Clinical therapeutics*, 30(4), 587-604.
- Clarke, P., Gray, A., & Holman, R. (2002). Estimating utility values for health states of type 2 diabetic patients using the EQ-5D (UKPDS 62). *Medical Decision Making*, 22(4), 340-349.