

- Marre, M., Shaw, J., Brändle, M., Bebakar, W. W., Kamaruddin, N. A., Strand, J., & Colagiuri, S. (2009). Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with Type 2 diabetes (LEAD-1 SU). *Diabetic Medicine*, 26(3), 268-278.
- Martínez, Y. V., Prado-Aguilar, C. A., Rascón-Pacheco, R. A., & Valdivia-Martínez, J. J. (2008). Quality of life associated with treatment adherence in patients with type 2 diabetes: a cross-sectional study. *BMC health services research*, 8(1), 164.
- McKeage, K., & Siddiqui, M. A. A. (2008). Amlodipine/Atorvastatin Fixed-Dose Combination. *American journal of cardiovascular drugs*, 8(1), 51-67.
- Meier, J. J. (2012). GLP-1 receptor agonists for individualized treatment of type 2 diabetes mellitus. *Nature Reviews Endocrinology*, 8(12), 728-742.
- Messerli, F. H., Bakris, G. L., Ferrera, R. D., Houston, M. C., Petrella, R. J., Flack, J. M., & Neutel, J. M. (2006). Efficacy and safety of coadministered amlodipine and atorvastatin in patients with hypertension and dyslipidemia: results of the AVALON trial. *The Journal of Clinical Hypertension*, 8(8), 571-583.
- Monesi, L., Baviera, M., Marzona, I., Avanzini, F., Monesi, G., Nobili, A., & Roncaglioni, M. C. (2012). Prevalence, incidence and mortality of diagnosed diabetes: evidence from an Italian population-based study. *Diabetic Medicine*, 29(3), 385-392.