

Nava-Ocampo AA, Velázquez-Armenta EY, Han JY, Koren G. Use of proton pump inhibitors during pregnancy and breastfeeding. *Can Fam Physician*. 2006; 52: 853-4 .

Nichols DJ, Muirhead GJ, Harness JA. Pharmacokinetics of sildenafil after single oral doses in healthy male subjects: absolute bioavailability, food effects and dose proportionality. *Br J Clin Pharmacol* 2002; 53 (Suppl. 1): 5S-12S.

Onal A.; Oztunç A. Development and validation of high performance liquid chromatographic method for the determination of esomeprazole in tablets, *J. Food Drug Anal.*, Vol. 14, No. 1, 2006, pp 12-18.

Patel, B.H.; Suhagia, B.N.; Patel, M.M.; Patel, J.R. Determination of pantoprazole, domperidone, rabeprazole, esomeprazole and itopride in pharmaceutical products by reverse phase liquid chromatography using single mobile phase, *Chromatographia*, 65(2007)11-12, pp-743-748.

Patil Shamkant S, Dhabale Pandurang N and Kuchekar Bhanudas S. Development and Statistical Validation of Spectrophotometric Method for Estimation of Esomeprazole in Tablet Dosage Form. *Asian J. Research Chem.* 2(2): April.-June, 2009.

Patterson B, Bedding A, Jewell H et al. The effect of intrinsic and extrinsic factors on the pharmacokinetic properties of tadalafil (IC351). *Int J Impot Res* 2001;.