

Kamiyama, E., Nakai, D., Mikkaichi, T., Okudaira, N., Okazaki, O., 2010. Interaction of angiotensin II type 1 receptor blockers with Pgp substrates in Caco-2 cells and hMDR1-expressing membranes. *Life Science*. 86 (1–2), 52–58.

Kaplan NM. Other forms of secondary hypertension. In: Kaplan NM, Lieberman E, eds. *Clinical hypertension*. 7th ed. Baltimore: Williams & Wilkins, 1998:395–406.

Karlova, M., Treichel, U., Malago, M., Frilling, A., Gerken, G., & Broelsch, C. E. (2000). Interaction of *Hypericum perforatum* (St. John's wort) with cyclosporin A metabolism in a patient after liver transplantation. *Journal of Hepatology*, 33(5), 853–855.

Karlson, B. W., Zetterstrand, S., Olofsson, B., & Elmfeldt, D. (2009). A dose-response analysis of candesartan-hydrochlorothiazide combination therapy in patients with hypertension. *Blood pressure*, 18(3), 149-156.

Kazakevich Y. and Lobrutto R. 2007. *HPLC for pharmaceutical scientists*. Wiley-Interscience. ISBN: 0470087943, 9780470087947.

Kim, N. D., Mehta, R., Yu, W., Neeman, I., Livney, T., Amichay, A., ... & Lansky, E. (2002). Chemopreventive and adjuvant therapeutic potential of pomegranate (*Punica granatum*) for human breast cancer. *Breast cancer research and treatment*, 71(3), 203-217.

Kirk, J. K. (1999). Angiotensin-II receptor antagonists: their place in therapy. *American family physician*, 59(11), 3140.

Kitahara, K., & Kawai, S. (2007). Cyclosporine and tacrolimus for the treatment of rheumatoid arthritis. *Current Opinion in Rheumatology*, 19(3), 238–245.

Kjeldsen, S. E., Jamerson, K. A., Bakris, G. L., Pitt, B., Dahlöf, B., Velazquez, E. J., ... & Weber, M. (2008). Predictors of blood pressure response to intensified and fixed combination treatment of hypertension: the ACCOMPLISH study. *Blood pressure*, 17(1), 7-17.

KK Pradhan, US Mishra, S Pattnaik, CK Panda, KC Sahu, (2011), Development and validation of a stability-indicating UV spectroscopic method for candesartan in bulk and formulations, *Indian Journal of Pharmaceutical Sciences*, 73(6), 693-696.

Kronbach, T., Fischer, V., & Meyer, U. A. (1988). Cyclosporine metabolism in human liver: identification of a cytochrome P-450III gene family as the major cyclosporine-metabolizing enzyme explains interactions of cyclosporine with other drugs. *Clinical Pharmacology and Therapeutics*, 43(6), 630–635.

Kubo, K., Kohara, Y., Imamiya, E., Sugiura, Y., Inada, Y., Furukawa, Y., ... & Naka, T. (1993). Nonpeptide angiotensin II receptor antagonists. Synthesis and biological activity of benzimidazolecarboxylic acids. *Journal of medicinal chemistry*, 36(15), 2182-2195.

Kwok, Y. C., Hsieh, D. P. H., & Wong, P. K. (2005). Toxicity identification evaluation (TIE) of pore water of contaminated marine sediments collected from Hong Kong waters. *Marine pollution bulletin*, 51(8), 1085-1091.