## List of Figures

Figure legends	Page No.
(Figure 1.1): Insulin-Primary Structure.	18
( <b>Figure 1.2</b> ): The main administration routes for insulin (Duan & Mao, 2010)	21
( <b>Figure 1.3</b> ): Schematic diagrams illustrating the absorption, distribution and elimination of insulin following oral or subcutaneous (S.C.) administration (Sonaje et al., 2010).	24
(Figure 1.4): Chemical structure of Chitosan.	28
(Figure 1.5): In-vivo efficiency of orally delivered insulin and chitosan/insulin NPs (Mukhopadhyay et al., 2013).	31
(Figure 1.6): Chemical structure of DAG.	33
(Figure 1.7): A model for cellular uptake and degradation of insulin (Duckworth et al., 1998).	39
( <b>Figure 2.1</b> ): Some of in situ perfusion instruments, (A): Low flow rate peristaltic pump, (B): SomnoSuite <sup>™</sup> Small Animal Anesthesia System, (C): Oxy-life oxygen concentrator, (D): Bio Rad 2110 Fraction Collector.	47
(Figure 2.2): The eversion of small intestine on a stainless steel rod then filling the everted intestine with the oxygenated Krebs buffer solution. Moreover the filled everted intestine was divided into sacs.	57
(Figure 2.3): Cannulation of exposed intestine at both ends then the isolated intestine was perfused in circulating system.	59
(Figure 2.4): Ligation of the portal vein and the inferior vena cava (IVC) beside the left kidney.	60
(Figure 2.5): Cannulation the portal vein with 16 GA catheter and perfusion with Krebs buffer, then cut the lower inferior vena cava.	61