

(Figure 3.12): Plot showing percentage of blood glucose levels of diabetic rats after in situ intestinal perfusion of Rh-insulin solution and placebo with SC insulin. (Data is as mean \pm SEM).	84
(Figure 3.13): Comparison percentage of blood glucose levels between normal and diabetic rats after in situ Rh-insulin solution perfusion. (Data is as mean \pm SEM).	85
(Figure 3.14): Comparison percentage of blood glucose levels between normal and diabetic rats after in situ placebo perfusion with SC insulin injection (1IU/ml). (Date is as mean \pm SEM).	85
(Figure 3.15): Plot showing percentage of blood glucose levels of normal rats after in situ intestinal perfusion of oral insulin-loaded nanoparticles formula and oily placebo with SC insulin. (Data is as mean \pm SEM).	86
(Figure 3.16): Plot showing percentage of blood glucose levels of diabetic rats after in situ intestinal perfusion of oral insulin-loaded nanoparticles formula and oily placebo with SC insulin. (Data is as mean \pm SEM).	86
(Figure 3.17): Comparison percentage of blood glucose levels between normal and diabetic rats after in situ oral insulin formula perfusion. (Data is as mean \pm SEM).	87
(Figure 3.18): Comparison percentage of blood glucose levels between normal and diabetic rats after in situ oily placebo perfusion with SC insulin injection (1IU/ml). (Date is as mean \pm SEM).	87
(Figure 3.19): Percentage of glucose level in normal rats after perfusion of oral insulin preparation and compared with perfusion of Rh-insulin solution (Date is as mean \pm SEM).	88
(Figure 3.20): Percentage of glucose level in normal rats after perfusion of oral insulin preparation and compared with perfusion of Rh-insulin solution (Date is as mean \pm SEM).	88
(Figure 3.21): Semilogarithmic plot showing the insulin level (μ IU/ml) obtained after bolus injection of insulin into the portal vein. Data are as mean of 8 rats \pm SEM.	89
(Figure 3.22): Percentage of glucose levels of normal rats after insulin perfusion in normal liver at different flow rates followed by SC injection (Data are as mean of 8 rats \pm SEM).	91
(Figure 3.23): Percentage of glucose levels of normal rats after insulin perfusion in diabetic liver at different flow rates followed by SC injection (Data are as mean of 8 rats \pm SEM).	91
(Figure 3.24): Comparison the percentage of blood glucose level of normal rats after insulin perfusion at 5 ml/min in normal and diabetic livers followed by SC insulin injection (Data are as mean of 8 rats \pm SEM).	92
(Figure 3.25): Comparison the percentage of blood glucose level of normal rats after insulin perfusion at 1 ml/min in normal and diabetic livers followed by SC insulin injection (Data are as mean of 8 rats \pm SEM).	92