

| | | |
|-----|---|-----|
| | interval) serum candesartan drug concentration after 4 hours of drug administration between single and combined drug use | |
| 26. | Diagram with error bars comparing the mean (with its 95% confidence interval) serum candesartan drug concentration after 6 hours of drug administration between single and combined drug use | 113 |
| 27. | Diagram with error bars comparing the mean (with its 95% confidence interval) serum candesartan drug concentration after half an hour of drug administration between single and combined drug use | 114 |
| 28. | Diagram with error bars comparing the mean (with its 95% confidence interval) serum candesartan drug concentration after one hour of drug administration between single and combined drug use | 115 |
| 29. | Diagram with error bars comparing the mean (with its 95% confidence interval) serum candesartan drug concentration after 2 hours of drug administration between single and combined drug use | 116 |
| 30. | Diagram with error bars comparing the mean (with its 95% confidence interval) serum candesartan drug concentration after 3 hours of drug administration between single and combined drug use | 117 |
| 31. | Diagram with error bars comparing the mean (with its 95% confidence interval) serum candesartan drug concentration after 4 hours of drug administration between single and combined drug use | 118 |
| 32. | Diagram with error bars comparing the mean (with its 95% confidence interval) serum candesartan drug concentration after 6 hours of drug administration between single and combined drug use | 119 |
| 33. | Candesartan blank chromatogram | |
| 34. | Candesartan zero chromatogram | |
| 35. | Candesartan LLOQ chromatogram | |
| 36. | Candesartan QC Mid chromatogram | |