

<b>Figure 3.12</b> Bench-top stability test calibration curve of propranolol in serum.	103
<b>Figure 3.13</b> Autosampler stability test calibration curve of propranolol in serum.	104
<b>Figure 3.14</b> Propranolol concentrations versus mean area ratios for each concentration of standard points.	106
<b>Figure 3.15</b> A processed blank serum sample.	110
<b>Figure 3.16</b> A processed zero serum sample spiked with sildenafil; the internal standard of propranolol analysis.	110
<b>Figure 3.17</b> A processed lower limit of quantification (50 ng/ml) for serum sample spiked with sildenafil; the internal standard of propranolol analysis.	111
<b>Figure 3.18</b> A processed upper limit of quantification (3000 ng/ml) for serum sample spiked with sildenafil; the internal standard of propranolol analysis.	111
<b>Figure 3.19</b> <i>In vivo</i> serum concentration versus time curves of propranolol in rats after a single oral dose of 20 mg/kg of propranolol, propranolol with 100 and 200 mg/kg of glucosamine. Each data point represents the mean $\pm$ SEM (n=7).	112
<b>Figure 3.20</b> <i>In vivo</i> serum concentration versus time curves of propranolol in rats after a single oral dose of 20 mg/kg of propranolol, propranolol with 5 and 9 mg/kg of cimetidine and rifampin, respectively. Each data point represents the mean $\pm$ SEM (n=5).	113
<b>Figure 3.21</b> Observed versus SimCYP-predicted propranolol plasma concentrations of 20 mg/kg propranolol, propranolol with 100 mg/kg glucosamine and propranolol with 200 mg/kg glucosamine.	116
<b>Figure 3.22</b> Time dependence curves of propranolol serum concentration in rats using <i>in situ</i> single pass intestinal perfusion. Rats were administered propranolol alone 1 mg/ml, propranolol with glucosamine 10 mg/ml, propranolol with cimetidine 1 mg/ml, and	118