

2.5.5 Preparation of 2-methyl-1-[4-(pyrrolidin-1-yl)but-2-yn-1-yl]-2,3-dihydro-1H-indole (AZ-6)




Figure 27: 2-methyl-1-[4-(pyrrolidin-1-yl)but-2-yn-1-yl]-2,3-dihydro-1H-indole.

The titled compound was prepared following the general procedure for synthesis of 2-methyl-1-[4-(amino-1-yl)but-2-yn-1-yl]-2,3-dihydro-1H-indole, AZ2-AZ7, yielded 2.3 g 90.4 %. **IR (NaCl, Cm^{-1}):** 2963 (ArH, stretch), 1607, 1460, 1431 (Ar,C=C, stretch), 1234, 1147 (Ar, C=C, bending), 749, 718 (ArH, bending). **$^1\text{H-NMR}$ (DMSO- d_6):** δ 1.22 (d, 3H, N-CH- CH_3), 1.89, 1.98, 2.72, 3.15 (m, various protons of cyclicamine), 3.06 (d, 1H, CH_2 -CH-N), 3.49, 3.89 (d, 2H, $J = 2.4$ Hz, CH_2 -C) due to long range coupling, 3.80 (m, 1H, $J = 6.15$ Hz, N- CH - CH_3), 3.73, 4.13 (d, 2H, $J = 2.4$ Hz, CH_2 -N) due to long range coupling, , 6.81-7.28 (m, 4H, ArH).