

## List of Figures

Figure No.	Caption	Page No.
1	Various effects of EGF receptor activation by Mitogen and other kinases	5
2	N-(3-ethynylphenyl)-6,7-bis (2-methoxyethoxy)-4-quinazolineamine	6
3	N-(3-Chloro-4-fluoro-phenyl) -7- methoxy -6- (3-morpholin-4-ylpropoxy) quinazoline -4-amine	7
4	N-(chloro-6-methylphenyl) -2- {{6-{{4-(2-hydroxyethyl)-1-piperazinyl}}-2-methyl-4-pyrimidinyl} amino-thiazol carboxamide	8
5	4-{{(4- methylpiperazin-1-yl) methyl}- N- {4-methyl- 3-{{(4- pyridine3 ylpyrimidin-2-yl) amino}}phenyl}}benzamide	9
6	4-(4-{{3-(4-chloro-3-(trifluoromethyl) phenyl} ureido} phenoxy)-N2-methylpyridine -2- carboxamide -4- methylbenzene sulfonate	9
7	N- (4-bromo-2-fluorophenyl) -6-methoxy-7 {{(1-methylpiperidin-4-yl) methoxy} quinazoline-4-amine	10
8	18,21- Dihydro-17-demethoxy-18,21-dideoxo-17-(2 propenylamino) geldanamycin	11
9	N- (2-diethylaminoethyl) -5- {{(Z)-(5-fluoro-2-oxo-1H-indol-3-ylidene) methyl} -2,4-dimethyl-1H-pyrrole-3-carboxamide	12
10	(1S, 4S, 7Z, 10S, 16E, 21R)-7-ethylidene-4,21-diisopropyl-2-oxa-12,13-dithia-5,8,20,23-Tetrazabicyclo{8,7,6}tricos-16-ene-3,6,9,19,22-pentone	13
11	Illustration of different steps in angiogenesis process	17
12	Different strategies for targeting angiogenesis inhibition	22
13	(RS)-2-(2,6-dioxopiperidin-3-yl)-1H-isoindole-1,3(2H)-dione	27
14	3-(4-amino-1-oxo 1,3-dihydro-2H-isoindol-2-yl) piperidine -2,6- dione	28
15	4-hydroxy-N,1-dimethyl-2-oxo-N- phenyl-1,2-dihydroquinoline-3-carboxamide	29
16	(2-{{(R)-2-methylpyrrolidine-2yl} -1H-benzimidazole-4-carboxamide)	29
17	Similarity in steric and electronic environment between Lenalidomide, Linomide and Aminoacetylenic derivatives of 2-methyindoline	30