

- Shepherd, J., Blauw, G. J., Murphy, M. B., Bollen, E. L., Buckley, B. M., Cobbe, S. M., & Westendorp, R. G. (2002). Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *The Lancet*, 360(9346), 1623-1630.
- Skyler, J. S., Bergenstal, R., Bonow, R. O., Buse, J., DEED-WANIA, P., Gale, E. A., & SHERWIN, R. (2009). American Diabetes Association American College of Cardiology Foundation American Heart Association.
- Smith, D. H. (2008). Comparison of angiotensin II type 1 receptor antagonists in the treatment of essential hypertension. *Drugs*, 68(9), 1207-1225.
- Sullivan, S. D., Alfonso-Cristancho, R., Conner, C., Hammer, M., & Blonde, L. (2009). Long-term outcomes in patients with type 2 diabetes receiving glimepiride combined with liraglutide or rosiglitazone. *Cardiovasc Diabetol*, 8, 12.
- Testa, M. A., & Simonson, D. C. (1998). Health economic benefits and quality of life during improved glycemic control in patients with type 2 diabetes mellitus: a randomized, controlled, double-blind trial. *Jama*, 280(17), 1490-1496.
- Turner, R. C., Holman, R. R., Cull, C. A., Stratton, I. M., Matthews, D. R., Frighi, V., & Sutton, P. J. (1998). Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *lancet*, 352(9131), 837-853.