

- Vaidyanathan, J., Choe, S., & Sahajwalla, C. G. (2012). Type 2 diabetes in pediatrics and adults: thoughts from a clinical pharmacology perspective. *Journal of pharmaceutical sciences*, 101(5), 1659-1671.
- Venkitachalam, L., Wang, K., Porath, A., Corbalan, R., Hirsch, A. T., Cohen, D. J., & Magnuson, E. A. (2012). Global variation in the prevalence of elevated cholesterol in outpatients with established vascular disease or 3 cardiovascular risk factors according to national indices of economic development and health system performance. *Circulation*, 125(15), 1858-1869.
- Walsh, P. C., Marschke, P., Ricker, D., & Burnett, A. L. (2000). Patient-reported urinary continence and sexual function after anatomic radical prostatectomy. *Urology*, 55(1), 58-61.
- Wang, Y. R., Alexander, G. C., & Stafford, R. S. (2007). Outpatient hypertension treatment, treatment intensification, and control in Western Europe and the United States. *Archives of internal medicine*, 167(2), 141-147.
- Wilson, P. W., D'Agostino, R. B., Levy, D., Belanger, A. M., Silbershatz, H., & Kannel, W. B. (1998). Prediction of coronary heart disease using risk factor categories. *Circulation*, 97(18), 1837-1847.
- Wolf-Maier, K., Cooper, R. S., Banegas, J. R., Giampaoli, S., Hense, H. W., Joffres, M., & Vescio, F. (2003). Hypertension prevalence and blood pressure levels in 6 European countries, Canada, and the United States. *Jama*, 289(18), 2363-2369.