

Prevalence of Food Addiction during Covid-19 Pandemic in Amman, Jordan: A Cross Sectional Study

Ruba Mousa Musharbash^{1*}, Nour Amin Elshahry², Andrea McGrattan³, Samantha Al-Alam⁴, Lara Badawi⁴

¹ Clinical dietitian, Ruba Musharbash Center for Nutrition Counseling, Amman, Jordan.

² Nutrition Department, Faculty of Pharmacy and Medical Sciences, University of Petra, Amman, Jordan

³ Research Associate & Dietitian, Population Health Sciences Institute, Newcastle University, NE4 5PL. UK

⁴ Nutritionist, Ruba Musharbash Center for Nutrition Counseling, Amman, Jordan.

*Corresponding author: Ruba Mousa Musharbash, Clinical dietitian, Ruba Musharbash Center for Nutrition Counseling.

Received date: April 13, 2021; Accepted date: April 30, 2021; Published date: May 05, 2021

Citation: Ruba M Musharbash, Nour A Elshahry, Andrea McGrattan, Samantha Al-Alam, Lara Badawi, (2021). Prevalence of Food Addiction during Covid-19 Pandemic in Amman, Jordan: A Cross Sectional study. *J. Nutrition and Food Processing*, 4(4); DOI:10.31579/2637-8914/050

Copyright: © 2021, Ruba Mousa Musharbash. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Aim: Emerging evidence highlights the impact of stressful conditions on eating behaviours, particularly the consumption of foods high in salt, sugar and fat. The impact of the Coronavirus disease-19 (COVID-19) pandemic on food addiction is still un-estimated. This study aimed to examine the prevalence of food addiction among the Jordanian population during the quarantine.

Materials and methods: A Cross-sectional study has been conducted between the 27th of April and the 4th of June of 2020. An online survey of Yale food addiction scale version.2.0 (YFAS 2.0) was distributed. A convenience sample was collected through social media sites. Differences in socio-demographic characteristics were analysed using t-test and chi-square tests. Food addiction was classified based on the scoring scale. Determinants of food addiction were predicted by using multivariate logistic regression.

Results: The results revealed that the prevalence of food addiction was 21.5% among adults. Among food addiction participants, 76.4% were diagnosed as severely food addicted. More than 50% of the participants reported high consumption of foods rich in sugar, salt, and fat (48.2%, 51.8%, and 52.5%, respectively). Overweight and obese young adults were more likely to have food addiction compared to normal weight. The multinomial logistic regression model revealed that there was no determinants of food addiction among adults.

Conclusion: During the first months of the quarantine, the prevalence of food addiction was higher among obese and overweight adults compared to normal-weight adults.

Keywords: food addiction, obesity, quarantine, COVID-19; food craving; chi-square test; insulin; leptin; ghrelin; homeostasis; etiology

1. Introduction

Coronavirus (COVID-19) is a severe respiratory syndrome caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) [31]. It spread rapidly and cases increased worldwide [46]. The World Health Organization (WHO) declared COVID-19 as a pandemic in March 2020. On the 17th of March, Jordan declared the national defence law, where a complete lockdown was launched with strict rules on citizens' mobility, to mitigate the spread of the virus [27]. The lockdown and quarantine, due to COVID-19, affected people's behaviors, economics [38], social and general health status [6]. Staying at home with limitations of outdoor activities, and less mobility, caused an increase in boredom and stress [31]. Consequently, all these stressful events are well known to affect eating patterns, and people's choices of food [21].

Additional stress can encourage people to overeat, especially craving specific comfort and palatable foods [37]. This intense desire to consume specific foods was previously described as a food craving, which is a

multidimensional concept, including emotional, cognitive and physiological processes [43]. The concept of Food Addiction (FA) can be presented as the phase where foods, mainly rich in sugar, salt and fat are potentially addictive, and that specific forms of overeating may be viewed as addictive behaviour [36]. It is now used to explain the etiology and maintenance of some forms of obesity [29]. The Yale food addiction scale 2nd version (YFAS 2.0) is used to examine FA, it is a diagnostic self-report questionnaire of addiction eating behaviour [1].

To the best of the authors' knowledge, the effect of COVID-19 on FA has not been studied before. Quarantine may have affected the eating behaviours of Jordanians. Thus, this study aimed to examine the FA prevalence among adults in Jordan during the first few weeks of the quarantine.

2. Materials and methods

2.1 Study Design

A cross-sectional study was conducted among Jordanian adults -aged more than 18 years. Data were collected from the 27th of April until the 4th of June 2020 by using an online questionnaire. The respondents were chosen using convenience sampling (non-probability sampling method) due to the lack of a population sampling system [10]. The online questionnaire opened from April to the end of June 2020. A consent form was included on the first page and 'agree and continue' choice could be selected for those who agreed to participate in the survey.

The guidelines for strengthening the Reporting of Observational Studies in Epidemiology (STROBE-nut) were used to write this article [51].

2.2 Study Tool

In the current study, the questionnaire included two parts; the first part included socio-demographic. The second part included YFAS 2.0, which is an instrument that assesses addictive-like eating behaviours using the criteria for substance use [28, 30]. Valid and reliable Arabic version copy was published in previous work in Jordan [16].

The scale is composed of a 35 item self-reported questionnaire that applies eleven Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV) criteria [36,42]. The scale gives two scoring options, the YFAS symptoms count [34,41], which indicates the number of dependence symptoms that have been met and the dichotomous version [31,50]. The measures of items are scored using a scale count ranging from never (0) to every day (7) indicating the number of dependence symptoms [28]. The participants with FA are diagnosed when having at least three symptoms with clinically significant impairment or distress [26,45]. The scale finally will subdivide the participants into four groups according to the severity of FA: mild, moderate, severe-FA and non-FA [34,41].

2.3 Participants

Participants were asked to complete the questionnaire and answer the questions related to their eating behaviours during the quarantine period from the beginning of the lockdown in Jordan.

The inclusion criteria of this study were: (1) adults aged ≥ 18 years old; (2) currently living in Amman (capital city of Jordan) during the lockdown due to the COVID -19 pandemic. Submitted responses that did

not match the inclusion criteria were excluded. The study was approved by the Institutional Review Board and the ethical committee of Petra University, Amman, Jordan (Grant number: 2Q/3/2020).

2.4 Statistical Analysis

Descriptive statistics, including frequencies, percentages, and cross-tabulation were collated to find the relationship between FA and socio-demographic characteristics of the participants. FA was classified as mild, moderate, severe based on the scale scoring [41] for the participants who have FA during the study time. The rest of the analysis was performed on the FA participants. Cross-tabulation was used to assess the relationship between the symptoms of FA and the FA categories. Multinomial logistic regression was used to predict the socio-demographic characteristics that could be determinants for FA among adults in this study. The significant level (p-value) in this study is ≤ 0.05 . The data were analysed using IBM SPSS Statistics for Windows, Version 26.0 (IBM Corp., Armonk, NY, USA). BMI was measured using self-reported weight and height and calculated by dividing weight in kilogram by height in meters squared [8].

3. Results

3.1 Participants Characteristics

The total number of 718 participants has completed the survey, only 652 met the inclusion criteria. The majority of the participants were women (94.2%), mean age was (30 \pm 7.7), and more than a quarter of the respondents were between 18-28 years old (46.62%). Moreover, the anthropometric results indicated that more than the half of participants were overweight or obese (59.6%), mean body mass index (BMI) was (26.9 \pm 5.5). Nearly half of participants consumed more foods that are rich in sugar, salt and fat occasionally (48.2%, 51.8%, 52.5%), respectively.

3.2 Prevalence of FA categories among the study participants

From the total sample, 21.5% (n=140) met the clinical threshold for FA. Based on YFAS scale score analysis, the FA is classified into three categories: mild, moderate and severe. Within the FA categories, the participants were distributed to 10%, 3.6% and 76.4% as mild-FA, moderate-FA and severe-FA as shown in Table 1.

Independent variables		N (%)	Non-FA n=512 n(%)	FA n=140 n (%)	P-Value
Age (years)	18-28	304 (46.6)	254 (49.6)	50 (35.7)	0.001
	29-39	278 (42.6)	199 (38.9)	79 (56.4)	
	40-50	58 (9)	47 (9.2)	11 (7.9)	
	51-60	12 (1.8)	12 (2.3)	0 (0)	
Gender	Women	614 (94.2)	481 (93.9)	133 (95)	0.64
	Men	38 (5.8)	31 (6.1)	7 (5)	
BMI (kg/m ²)	Underweight	13(2)	12 (2.3)	1 (0.7)	<0.001
	Normal weight	250 (38.3)	228 (44.5)	22 (15.7)	
	Overweight	239 (36.7)	176 (34.4)	63 (45)	
	Obese	150 (23)	96 (18.8)	54 (38.6)	
Salary (JOD)	Less than 300	67 (10.3)	50 (9.8)	17 (12.1)	0.79
	Between 300-500	154 (23.6)	119 (23.2)	35 (25.0)	
	Between 500-700	207 (31.7)	164 (32.0)	43 (30.7)	
	more than 1000	224 (34.4)	179 (35.0)	45 (32.2)	
Education level	High school or less	91 (14)	67 (13.1)	24 (17.1)	0.34
	Bachelors or diploma	496 (76.1)	396 (77.3)	100 (71.4)	
	Higher education	65 (9.9)	49 (9.6)	16 (11.4)	
Smoker	Yes	87 (13.3)	63 (12.3)	24 (17.1)	0.14
	No	565 (86.7)	449 (87.7)	116 (82.9)	
Frequency of consuming foods high in sugar	Don't consume them	12 (1.8)	9 (1.8)	3 (2.1)	<0.001
	Rarely	108 (16.6)	101 (19.7)	7 (5.0)	

	Occasionally	314 (48.2)	259 (50.6)	55 (39.3)	
	Daily	218 (33.4)	143 (27.9)	75 (53.6)	
Frequency of consuming foods high in salt	Don't consume them	36 (5.52)	29 (5.7)	7 (5.0)	<0.001
	Rarely	141 (21.63)	127 (24.8)	14 (10)	
	Occasionally	338 (51.84)	263 (51.4)	75 (53.6)	
	Daily	137 (21.01)	93 (18.2)	44 (31.4)	
Frequency of consuming foods high in fat	Don't consume them	44 (6.7)	40 (7.8)	4 (2.9)	<0.001
	Rarely	215 (33)	188 (36.7)	27 (19.3)	
	Occasionally	342 (52.5)	257 (50.2)	85 (60.7)	
	Daily	51 (7.8)	27 (5.3)	24 (17.1)	
P-value ≤0.05. BMI, body mass index; JOD, Jordanian Dinar; FA, food addiction; Non-FA, non-food addiction					

Table 1. Socio-demographic characteristics of participants according to the food addiction groups (FA and Non-FA), n=652

3.3 Relationship between FA groups and the socio-demographic characteristics of the study population

As shown in Table 2, The relation between age, BMI, frequency of eating foods rich in (salt, sugar and fat) and FA variables were significant, $X^2(1, N = 642) = 16.002, p= 0.001, = 47.434, p= 0.001, = 20.539, p= 0.001, = 39.016, p= 0.001, = 36.253, p= 0.001$, respectively. Adults who are 29-39 years old, overweight, eating foods high in sugar daily and occasionally eating foods high in salt and fat were more likely to be food addicted than others.

3.4 Relationship between FA categories (mild, moderate and severe) and the socio-demographic characteristics of the study population

As shown in Table 2, a chi-square test of independence was performed to examine the relationship between socio-demographic and eating lifestyle and the FA categories. The relation between frequency of consuming foods rich in sugar and FA categories was significant, $X^2(1, N = 642) = 19.28, p= 0.004$.

Independent variables		Mild-FA n=14 n (%)	Moderate-FA n=19 n (%)	Severe-FA n=107 n (%)	P-Value
Age (years)	18-28	4 (8)	6 (12)	40 (80)	0.49
	29-39	8 (10.1)	10 (12.7)	61 (77.2)	
	40-50	2 (18.2)	3 (27.3)	6 (54.5)	
	51-60	0 (0)	0 (0)	0 (0)	
Gender	Women	14 (10.5)	17 (12.8)	102 (76.7)	0.37
	Men	0 (0)	2 (28.6)	5 (71.4)	
BMI (kg/m ²)	Underweight	0 (0)	1 (100)	0 (0)	0.19
	Normal weight	4 (18.2)	3 (13.6)	15 (68.2)	
	Overweight	5 (7.9)	9 (14.3)	49 (77.8)	
	Obese	5 (9.3)	6 (11.1)	43 (79.6)	
Salary (JOD)	Less than 300	2 (11.8)	3 (17.6)	12 (70.6)	0.98
	Between 300-500	3 (8.6)	6 (17.1)	26 (74.3)	
	Between 500-700	4 (9.3)	5 (11.6)	34 (79.1)	
	more than 1000	5 (11.1)	5 (11.1)	35 (77.8)	
Education level	High school or less	4 (16.7)	2 (8.3)	18 (75)	0.64
	Bachelors or diploma	8 (8)	14 (14)	78 (78)	
	Higher education	2 (12.5)	3 (18.7)	11 (68.8)	
Smoker	Yes	4 (16.7)	4 (16.7)	16 (66.7)	0.39
	No	10 (8.6)	15 (13)	91 (78.4)	
P-value <0.05. BMI, body mass index; JOD, Jordanian Dinar; FA: food addiction; Non-FA: non-food addiction.					

Table 2. Socio-demographic characteristics of participants according to the food addiction categories, n=140

Adults who eat foods high in sugar are more likely to be severely food addicted than others.

prediction between moderate-FA and consumption of frequency of consuming foods high in sugar ($B=-0.993, s.e.= .379, p 0.014$) and these results are consistent with the chi-square test.

In Table 3 the multinomial logistic regression results present a significant

Variables	Mild-FA		Moderate-FA	
	B	Exp(B) – (95% CI)	B	Exp(B) – (95% CI)
BMI	-0.40	0.67 (0.29 - 1.52)	-0.61	0.54 (.26 - 1.13)
Gender	-17.41	2.74 (2.74 - 2.74)	2.00	7.10 (0.80 - 63.51)
Salary (JOD)	-0.08	0.93 (0.48 - 1.81)	-0.50	0.61 (0.33 - 1.13)

Education	-0.29	0.75 (0.23 - 2.44)	0.70	2.00 (0.63 - 6.04)
Smoker	-0.96	0.38 (0.10 - 1.54)	-0.36	0.70 (0.17 - 2.92)
Frequency of consuming foods high in sugar	-0.81	0.44 (0.20 - 1.01)	-0.93	0.40 (0.19 - 0.83)
Frequency of eating salty foods	-0.33	0.72 (0.33- 1.60)	-0.29	0.80 (0.37 - 1.52)
Frequency of eating high fat food	-0.66	0.52 (0.21 - 1.27)	-0.41	0.66 (0.29 - 1.52)
The reference category is: Sever-FA. P-value ≤ 0.05 . BMI, body mass index; JOD, Jordanian Dinar; FA, food addiction; Non-FA, non-food addiction				

Table 3. Multivariate logistic regression of the Socio-demographic characteristics and FA categories, n=140

3.5 The relationship between FA symptom scores and frequency of symptoms among food addicted participants.

The mean symptoms score for the whole sample was 2.95 ± 3.50 , and 7.95 ± 2.85 for the FA group, Table 4. Illustrates the percentages of individuals with different levels of FA that reported each of the eleven symptoms. The most frequent symptoms reported in the whole sample were "Important social, occupational, or recreational activities given up or reduced" (36.5%), and "Characteristic withdrawal symptoms;

substance is taken to relieve withdrawal" (34.7%). Among individuals who have severe FA the main symptoms reported were, "Characteristic withdrawal symptoms; substance taken to relieve withdrawal" (82.1%), followed by "Persistent desire or repeated unsuccessful attempts at quitting symptom" (84.4%). Meanwhile, the less frequent symptom in both the whole sample and in the group, who met the FA criteria was "Use continues despite knowledge of adverse consequences (e.g., failure to fulfil role obligation, use when physically hazardous)" 16.7%, 60%, respectively.

Symptoms	Mild-FA n=14 n (%)	Moderate-FA n=19 n (%)	Severe-FA n=107 n (%)	P-Value
Substance taken in larger amount and for longer period than intended	7 (6.7)	6 (5.7)	92 (87.6)	<0.001
Persistent desire or repeated unsuccessful attempts at quitting	7 (5.7)	12 (9.8)	103 (84.4)	<0.001
Much time/activity to obtain, use, recover	2 (2)	11 (11)	87 (87)	<0.001
Important social, occupational, or recreational activities given up or reduced	3 (2.7)	9 (8.1)	99 (89.2)	<0.001
Use continues despite knowledge of adverse consequences (e.g., failure to fulfil role obligation, use when physically hazardous	1 (1.2)	4 (4.8)	79 (94)	<0.001
Tolerance (marked increase in amount; marked decrease in effect)	0 (0)	6 (6.5)	87 (93.5)	<0.001
Characteristic withdrawal symptoms; substance taken to relieve withdrawal	7 (5.7)	15 (12.2)	101 (82.1)	<0.001
Continued use despite social or interpersonal problems	1 (1.1)	5 (5.5)	85 (93.4)	<0.001
Failure to fulfil major role obligation (e.g., work, school, home)	1 (1.2)	3 (3.5)	81 (95.3)	<0.001
Use in physically hazardous situations	3 (3.2)	5 (5.4)	85 (91.4)	<0.001
Craving, or a strong desire or urge to use	2 (1.9)	10 (9.4)	94 (88.7)	<0.001
Data are presented as numbers (percentages). P-value ≤ 0.05				

Table 4. Food addiction symptom scores and frequency of symptoms in participants with Food addiction, n=140.

Discussion

Many studies have been published recently on eating behaviours during the COVID-19 quarantine [12,20,21]. To our knowledge, no studies are available on the prevalence of FA during the COVID-19 pandemic quarantine.

The percentage of participants who are food addicted in our study is considered high compared to other countries [4,5,45]. A previous study conducted in Jordan concluded that the prevalence of FA (n=4479) was 36.7%, FA was significantly greater among young adults aged 18-34 years, overweight and obese participants [16]. Another study in Egypt

concluded that the FA prevalence was 15.7% in the studied sample [26]. Furthermore, a cohort study additionally reported that FA among middle aged US young women was higher among the younger proportion, and the majority of the population that was diagnosed with FA fell into the overweight category [9]. Obesity and FA have been the most studied in the literature; recent meta-analysis studies concluded that the prevalence of FA was higher in the population that fell in the overweight and obese persons [36,39].

However, Jordan is considered one of the highest countries in the prevalence of obesity and overweight, consisting of 69.6% of the whole population who are overweight [52]. Most of the women with FA in our

sample were overweight or obese, and this was also evidenced in previous studies [7,17,18,19,24]. Results from the same studies showed that youth scored the most for severe FA [45,36]. One possible explanation could be that respondents from young age groups are more exposed to food cues and changes in their dietary lifestyle and habits. Furthermore, they may be at greater risk of emotion-induced eating and exposure to stress factors [36]. The stress during the quarantine could be related to changing eating behaviours and emotional eating during the lockdown time. A very recent study in Saudi Arabia concluded that half of the women in the study sample have suffered from emotional eating during quarantine, and increased intake of foods rich in sugar and fat markedly with increased stress levels [2]. A similar study during quarantine reported an increase of daily food intake by almost 50% of the respondents and by the variety of food choices; hence, consuming fewer fruits and vegetables and consuming dairy, meat and fast foods [12,21].

Stress can disrupt eating behaviour and encourage eating high palatable foods [14]. Therefore, the activation of the reward circuit; hence the mesolimbic dopaminergic system secretes dopamine in the ventral tegmental area (VTA) [16,25]. Insulin, leptin, and ghrelin also play a key role in the homeostasis of eating behaviors [11,23]. Leptin decreases food intake and decreases the firing rate of dopamine release through signalling in the ventral tegmental area (VTA) [15]. Ghrelin, leptin antagonist, increases food intake and influences the signaling in the VTA [22]. Insulin increases food intake by increasing the release and firing of dopamine in the VTA [3,13].

The results of the current study showed that some of the respondents had consumed palatable foods during the stressful condition of quarantine due to the COVID-19 pandemic. Some studies were in agreement as they mentioned that the pandemic changed the consumption patterns [32]. Respondents reported high-stress levels during the COVID-19 pandemic that led to the increased binge-eating [48]. Stress and emotional eating have also been studied as there was a strong positive relationship between increased emotional eating and stress induced by quarantine [47,53].

The mean symptoms score in the whole sample was high compared with other studies [1,29,49]. Also, higher in FA group, compared to in a previous Jordanian study [16].

There were many limitations to consider in this study. First, this study does not compare the presence of FA before and after the pandemic. Therefore, the results compared with different populations. Second, the majority of the participants who completed the online survey were women, this may be explained by higher interest in health topics in women compared to men. The strengths of this study are that it is the first study globally and in Jordan discussing the FA status during the quarantine, and the use of a strong scale to measure FA among the participants. More longitudinal studies are required in the future.

4. Conclusion

In summary, the prevalence of FA and the mean symptoms score were high in the FA group compared to other previous studies. Younger aged adults and overweight respondents were more likely to have FA. Intervention studies are required to manage eating behaviours among adults during times of increased stress. This study recommends the need for more longitudinal studies on FA to coincide with the increasing quarantine duration in Jordan and other countries. Moreover, awareness training and programs are recommended to increase the responsiveness toward eating and stress management and its short and long term health effects among adults.

References

- Ahmad, M., & Musharbash, R. (2018). Examining the Connection between Food Addiction and Human Obesity. *Nature and Science*, 16, 102–110.
- Al-Musharaf, S. (2020). Prevalence and Predictors of Emotional Eating among Healthy Young Saudi Women during the COVID-19 Pandemic. *Nutrients*, 12(10), 2923.
- Bruijnzeel, A. W., Corrie, L. W., Rogers, J. A., & Yamada, H. (2011). Effects of insulin and leptin in the ventral tegmental area and arcuate hypothalamic nucleus on food intake and brain reward function in female rats. *Behavioural Brain Research*, 219(2), 254–264.
- Burrows, T., Hides, L., Brown, R., Dayas, C., & Kay-Lambkin, F. (2017). Differences in Dietary Preferences, Personality and Mental Health in Australian Adults with and without Food Addiction. *Nutrients*, 9(3), 285.
- Ceccarini, M., Manzoni, G. M., Castelnuovo, G., & Molinari, E. (2015). An Evaluation of the Italian Version of the Yale Food Addiction Scale in Obese Adult Inpatients Engaged in a 1-Month-Weight-Loss Treatment. *Journal of Medicinal Food*, 18(11), 1281–1287.
- de Lima, C. V. C., Cândido, E. L., da Silva, J. A., Albuquerque, L. V., Soares, L. de M., do Nascimento, M. M., de Oliveira, S. A., & Neto, M. L. R. (2020). Effects of quarantine on mental health of populations affected by Covid-19. *Journal of Affective Disorders*, 275, 253–254.
- Erzsébet Magyar, É., Tényi, D., Gearhardt, A., Jeges, S., Abaliget, G., Tóth, Á. L., Janszky, J., Molnár, D., Tényi, T., & Csábi, G. (2018). Adaptation and validation of the Hungarian version of the Yale Food Addiction Scale for Children. *Journal of Behavioral Addictions*, 7(1), 181–188.
- Flegal, K.M. Carroll, M.D. Kit, B.K. and Ogden, C. L. (2012), Prevalence of obesity and trends in the distribution of body mass index among adults, 1999-2010. *American Medical Association*, 307 (5): 491-497.
- Flint, A. J., Gearhardt, A. N., Corbin, W. R., Brownell, K. D., Field, A. E., & Rimm, E. B. (2014). Food-addiction scale measurement in 2 cohorts of middle-aged and older women. *The American Journal of Clinical Nutrition*, 99(3), 578–586.
- Hajian-Tilaki, K. (2011). Sample size estimation in epidemiologic studies. *Caspian Journal of Internal Medicine*, 2(4), 289–298.
- Keen-Rhinehart, E., Ondek, K., & Schneider, J. E. (2013). Neuroendocrine regulation of appetitive ingestive behavior. *Frontiers in Neuroscience*, 7, 213.
- Kriaucioniene, V., Bagdonaviciene, L., Rodríguez-Pérez, C., & Petkeviciene, J. (2020). Associations between Changes in Health Behaviours and Body Weight during the COVID-19 Quarantine in Lithuania: The Lithuanian COVIDiet Study. *Nutrients*, 12(10), 3119.
- Labouèbe, G., Liu, S., Dias, C., Zou, H., Wong, J. C. Y., Karunakaran, S., Cleo, S. M., Phillips, A. G., Boutrel, B., & Borgland, S. L. (2013). Insulin induces long-term depression of ventral tegmental area dopamine neurons via endocannabinoids. *Nature Neuroscience*, 16(3), 300–308.
- Lin, Y.-S., Tung, Y.-T., Yen, Y.-C., & Chien, Y.-W. (2020). Food Addiction Mediates the Relationship between Perceived Stress and Body Mass Index in Taiwan Young Adults. *Nutrients*, 12(7).
- Murray, S., Tulloch, A., Gold, M. S., & Avena, N. M. (2014). Hormonal and neural mechanisms of food reward, eating behaviour and obesity. *Nature Reviews. Endocrinology*, 10(9), 540–552.
- Musharbash, R., Ahmad, M. (2018). Food addiction: its prevalence and relation with insulin resistance, lipid profile, c-reactive protein and anthropometric indices, and effect of weight-

- reducing diet in a sample of obese young adults in amman [Unpublished doctoral thesis]. The University of Jordan
17. Nunes-Neto, P. R., Köhler, C. A., Schuch, F. B., Solmi, M., Quevedo, J., Maes, M., Murrú, A., Vieta, E., McIntyre, R. S., McElroy, S. L., Gearhardt, A. N., Stubbs, B., & Carvalho, A. F. (2018). Food addiction: Prevalence, psychopathological correlates and associations with quality of life in a large sample. *Journal of Psychiatric Research*, 96, 145–152.
 18. Pedram, P., Wadden, D., Amini, P., Gulliver, W., Randell, E., Cahill, F., Vasdev, S., Goodridge, A., Carter, J. C., Zhai, G., Ji, Y., & Sun, G. (2013). Food Addiction: Its Prevalence and Significant Association with Obesity in the General Population. *PLoS ONE*, 8(9), e74832.
 19. Pursey, K. M., Stanwell, P., Gearhardt, A. N., Collins, C. E., & Burrows, T. L. (2014). The prevalence of food addiction as assessed by the Yale Food Addiction Scale: a systematic review. *Nutrients*, 6(10), 4552–4590.
 20. Ruiz-Roso, M. B., de Carvalho Padilha, P., Mantilla-Escalante, D. C., Ulloa, N., Brun, P., Acevedo-Correa, D., Arantes Ferreira Peres, W., Martorell, M., Aires, M. T., de Oliveira Cardoso, L., Carrasco-Marín, F., Paternina-Sierra, K., Rodriguez-Meza, J. E., Montero, P. M., Bernabè, G., Pualetto, A., Taci, X., Visioli, F., & Dávalos, A. (2020). Covid-19 Confinement and Changes of Adolescent's Dietary Trends in Italy, Spain, Chile, Colombia and Brazil. *Nutrients*, 12(6), 1807.
 21. Sidor, A., & Rzymiski, P. (2020). Dietary Choices and Habits during COVID-19 Lockdown: Experience from Poland. *Nutrients*, 12(6), 1657.
 22. Skibicka, K. P., Hansson, C., Alvarez-Crespo, M., Friberg, P. A., & Dickson, S. L. (2011). Ghrelin directly targets the ventral tegmental area to increase food motivation. *Neuroscience*, 180, 129–137.
 23. Suzuki, K., Simpson, K. A., Minnion, J. S., Shillito, J. C., & Bloom, S. R. (2010). The role of gut hormones and the hypothalamus in appetite regulation. *Endocrine Journal*, 57(5), 359–372.
 24. Yang, F., Liu, A., Li, Y., Lai, Y., Wang, G., Sun, C., Sun, G., Shan, Z., & Teng, W. (2017). Food Addiction in Patients with Newly Diagnosed Type 2 Diabetes in Northeast China. *Frontiers in Endocrinology*, 8, 218.
 25. Yau, Y. H. C., & Potenza, M. N. (2013). Stress and eating behaviors. *Minerva Endocrinologica*, 38(3), 255–267.
 26. Ahmed, A. Y., & Sayed, A. M. (2017). Prevalence of food addiction and its relationship to body mass index. *Egyptian Journal of Medical Human Genetics*, 18(3), 257–260.
 27. Al-Tammemi, A. B. (2020). The Battle against COVID-19 in Jordan: An Early Overview of the Jordanian Experience. *Frontiers in Public Health*, 8.
 28. Chao, A. M., Grilo, C. M., & Sinha, R. (2016). Food cravings, binge eating, and eating disorder psychopathology: Exploring the moderating roles of gender and race. *Eating Behaviors*, 21, 41–47.
 29. Chao, A. M., Shaw, J. A., Pearl, R. L., Alamuddin, N., Hopkins, C. M., Bakizada, Z. M., Berkowitz, R. I., & Wadden, T. A. (2017). Prevalence and psychosocial correlates of food addiction in persons with obesity seeking weight reduction. *Comprehensive Psychiatry*, 73, 97–104.
 30. Davis, C., Curtis, C., Levitan, R. D., Carter, J. C., Kaplan, A. S., & Kennedy, J. L. (2011). Evidence that 'food addiction' is a valid phenotype of obesity. *Appetite*, 57(3), 711–717.
 31. Di Renzo, L., Gualtieri, P., Pivari, F., Soldati, L., Attinà, A., Cinelli, G., Leggeri, C., Caparello, G., Barrea, L., Scerbo, F., Esposito, E., & De Lorenzo, A. (2020). Eating habits and lifestyle changes during COVID-19 lockdown: An Italian survey. *Journal of Translational Medicine*, 18(1), 229.
 32. Eftimov, T., Popovski, G., Petković, M., Seljak, B. K., & Kocev, D. (2020). COVID-19 pandemic changes the food consumption patterns. *Trends in Food Science & Technology*, 104, 268–272.
 33. Gearhardt, A. N., Corbin, W. R., & Brownell, K. D. (2016). Development of the Yale Food Addiction Scale Version 2.0. *Psychology of Addictive Behaviors*, 30(1), 113–121.
 34. Gearhardt, A. N., Roberto, C. A., Seamans, M. J., Corbin, W. R., & Brownell, K. D. (2013). Preliminary validation of the Yale Food Addiction Scale for children. *Eating Behaviors*, 14(4), 508–512.
 35. Gearhardt, A., Roberts, M., & Ashe, M. (2013). If Sugar is Addictive... What Does It Mean for the Law? *The Journal of Law, Medicine & Ethics*, 41(1_suppl), 46–49.
 36. Hauck, C., Weiß, A., Schulte, E. M., Meule, A., & Ellrott, T. (2017). Prevalence of "Food Addiction" as Measured with the Yale Food Addiction Scale 2.0 in a Representative German Sample and Its Association with Sex, Age and Weight Categories. *Obesity Facts*, 10(1), 12–24.
 37. Jacques, A., Chaaya, N., Beecher, K., Ali, S. A., Belmer, A., & Bartlett, S. (2019). The impact of sugar consumption on stress driven, emotional and addictive behaviors. *Neuroscience & Biobehavioral Reviews*, 103, 178–199.
 38. Legese Feyisa, H. (2020). The World Economy at COVID-19 Quarantine: Contemporary Review. *International Journal of Economics, Finance and Management Sciences*, 8(2), 63.
 39. Lennerz, B., & Lennerz, J. K. (2018). Food addiction, high-glycemic-index carbohydrates, and obesity. *Clinical Chemistry*, 64(1), 64–71.
 40. MacDonald, P. L., & Gardner, R. C. (2000). Type I Error Rate Comparisons of Post Hoc Procedures for I j Chi-Square Tables. *Educational and Psychological Measurement*, 60(5), 735–754.
 41. Meule, A., Gearhardt, A. (2019). Ten Years of the Yale Food Addiction Scale: a Review of Version 2.0. *Current Addiction Reports*, 6, 218–228.
 42. Meule, A., Hermann, T., & Kübler, A. (2015). Food addiction in overweight and obese adolescents seeking weight-loss treatment. *European Eating Disorders Review*, 23(3), 193–198.
 43. Muscogiuri, G., Barrea, L., Savastano, S., & Colao, A. (2020). Nutritional recommendations for CoVID-19 quarantine. *European Journal of Clinical Nutrition*, 74(6), 850–851.
 44. Ouellette, A.-S., Rodrigue, C., Lemieux, S., Tchernof, A., Biertho, L., & Bégin, C. (2017). Yale Food Addiction Scale: Examining the Psychometric Properties of the French Version among Individuals with Severe Obesity Awaiting Bariatric Surgery. *Psychology*, 08(14), 2547–2561.
 45. Pursey, K. M., Collins, C. E., Stanwell, P., & Burrows, T. L. (2016). The stability of "food addiction" as assessed by the Yale Food Addiction Scale in a non-clinical population over 18-months. *Appetite*, 96, 533–538.
 46. Singhal, T. (2020). A Review of Coronavirus Disease-2019 (COVID-19). *The Indian Journal of Pediatrics*, 87(4), 281–286.
 47. Shen, W., Long, L. M., Shih, C.-H., & Ludy, M.-J. (2020). A Humanities-Based Explanation for the Effects of Emotional Eating and Perceived Stress on Food Choice Motives during the COVID-19 Pandemic. *Nutrients*, 12(9), 2712.
 48. Termorshuizen, J. D., Watson, H. J., Thornton, L. M., Borg, S., Flatt, R. E., MacDermod, C. M., Harper, L. E., Furth, E. F., Peat, C. M., & Bulik, C. M. (2020). Early impact of COVID-19 on individuals with self-reported eating disorders: A survey of ~1,000 individuals in the United States and the Netherlands. *International Journal of Eating Disorders*, eat.23353.
 49. Tompkins, C. L., Laurent, J., & Brock, D. W. (2017). Food Addiction: A Barrier for Effective *Weight Management for Obese Adolescents*. *Childhood Obesity*, 13(6), 462–469.

50. Torres, S., Camacho, M., Costa, P., Ribeiro, G., Santos, O., Vieira, F. M., Brandão, I., Sampaio, D., & Oliveira-Maia, A. J. (2017). Psychometric properties of the Portuguese version of the Yale Food Addiction Scale. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 22(2), 259–267.
51. Von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., Vandenbroucke, J. P., & STROBE Initiative (2014). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *International journal of surgery* (London, England), 12(12), 1495–1499.
52. WHO. (2016). Overweight and obesity.
53. Wiss, D. A., & Avena, N. M. (2020). Food Addiction, Binge Eating, and the Role of Dietary Restraint: Converging Evidence from Animal and Human Studies. In *Binge Eating* (pp. 193–209). Springer International Publishing.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Manuscript](#)

DOI: [10.31579/2637-8914/050](https://doi.org/10.31579/2637-8914/050)

Ready to submit your research? Choose Auctores and benefit from:

- ❖ fast, convenient online submission
- ❖ rigorous peer review by experienced research in your field
- ❖ rapid publication on acceptance
- ❖ authors retain copyrights
- ❖ unique DOI for all articles
- ❖ immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more www.auctoresonline.org/journals/nutrition-and-food-processing